

Quantum Field Theory with Null-Fronted Metrics

Orit Levin and Asher Peres

Department of Physics, Technion—Israel Institute of Technology, 32 000 Haifa, Israel

There is a large class of classical null-fronted metrics in which a free scalar field has an infinite number of conservation laws. In particular, if the scalar field is quantized, the number of particles is conserved. However, with more general null-fronted metrics, field quantization cannot be interpreted in terms of particle creation and annihilation operators, and the physical meaning of the theory becomes obscure.

1994 PACS numbers: 04.62.+v 04.20.Cv 03.50.-z

arXiv:hep-th/9401160v1 31 Jan 1994

Please address all correspondence to the second author.

E-mail: peres@techunix.technion.ac.il

Fax: + 972 4 221514

Conventional canonical dynamics, and likewise quantum field theory, use a time-like coordinate as the evolution parameter. However, it was pointed out long ago by Dirac [1] that it may be advantageous to use a null coordinate for that purpose. The mathematical consistency of quantum field theory on null slabs in Minkowski space was formally proved by Leutwyler, Klauder, and Streit [2]. This approach has led to numerous applications in particle physics [3].

In this work, we examine the consistency of canonical (or quantum) field theory when a null coordinate is used as the evolution parameter in curved spacetime. This choice may be advantageous for additional reasons: in some cases, a complete foliation of spacetime may be possible with null hypersurfaces, while it is impossible with space-like hypersurfaces. We start by giving a very simple derivation of a result which has been known for some time: classical plane-fronted gravitational waves do not polarize the vacuum; namely, their interaction with a free quantized field does not create particles [4–6]. We show that this property is a special case of an infinity of conservation laws, valid for any metric which obeys the conditions

$$g^{00} = g^{01} = g^{02} = 0, \quad (1)$$

and

$$\partial g^{\alpha\beta} / \partial x^3 = 0. \quad (2)$$

However, when we consider more general metrics, we run into severe consistency problems. In particular, a free quantum field cannot be interpreted as consisting of particle creation and annihilation operators.

The condition $g^{00} = 0$ means that hypersurfaces with constant $x^0 \equiv u$ (which will be used as “time” in the following discussion), have a null normal vector $\partial_\alpha u$. This is why u is called a “null coordinate.” Equation (2) is the definition of the “null-fronted” symmetry. We shall use numerical indices 0, 1, 2, 3, to denote the coordinates u, x, y, v , respectively. Greek indices refer to all four coordinates, and Latin indices to coordinates 1 and 2 only. Plane-fronted gravitational waves [7] are a special case of null-fronted metric: the only nonvanishing components of $g^{\alpha\beta}$ are $g^{03} = -g^{11} = -g^{22} = 1$, and g^{33} which satisfies $\sum_m \partial^2 g^{33} / (\partial x^m)^2 = 0$. Null-fronted metrics with more general values of $g^{\alpha\beta}$ indicate the presence of a nonvanishing energy-momentum tensor, which may sometimes (but not always) be physically realizable [8].

Equation (2), together with the geodesic equation,

$$\frac{d}{ds} \left(g_{\lambda\mu} \frac{dx^\mu}{ds} \right) = \frac{1}{2} \frac{\partial g_{\alpha\beta}}{\partial x^\lambda} \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds}, \quad (3)$$

which describes the motion of a free classical particle, implies that the momentum component,

$$p_3 \equiv m g_{3\alpha} (dx^\alpha / ds), \quad (4)$$

is a constant of the motion (m is the mass of the particle). Since *all* classical orbits have this property, we expect the existence of infinitely many conservation laws in a free field theory (classical or quantal) having a null-fronted background metric.

Consider indeed a free classical real scalar field ϕ , satisfying the Klein-Gordon equation $(\gamma^{\alpha\beta} \phi_{,\alpha})_{,\beta} = -\kappa^2 \gamma \phi$, where $\gamma \equiv (-\text{Det } g_{\alpha\beta})^{1/2}$ and $\gamma^{\alpha\beta} \equiv \gamma g^{\alpha\beta}$. (The $\gamma^{\alpha\beta}$ are denoted by

Gothic letters in the older literature.) All these expressions are functions of u , x , and y only. Explicitly, the Klein-Gordon equation is

$$2\gamma^{03}\phi_{,03} + \gamma^{03}_{,0}\phi_{,3} + \gamma^{33}\phi_{,33} + 2\gamma^{m3}\phi_{,m3} + \gamma^{m3}_{,m}\phi_{,3} + (\gamma^{mn}\phi_{,m})_{,n} = -\kappa^2\gamma\phi. \quad (5)$$

Let us perform a Fourier transformation with respect to v (only):

$$\chi(u, x, y, p) \equiv (2\pi)^{-1/2} \int_{-\infty}^{\infty} e^{ivp} \phi(u, x, y, v) dv. \quad (6)$$

For quantized particles, the physical meaning of the parameter p would be $p_3/\hbar$. However, the present argument is strictly classical (quantization will be discussed later). The Fourier transform of Eq. (5) is

$$-2ip\gamma^{03}\chi_{,0} - ip\gamma^{03}_{,0}\chi - p^2\gamma^{33}\chi - 2ip\gamma^{m3}\chi_{,m} - ip\gamma^{m3}_{,m}\chi + (\gamma^{mn}\chi_{,m})_{,n} = -\kappa^2\gamma\chi. \quad (7)$$

Let us further define $\psi \equiv \sqrt{\gamma^{03}}\chi$. After some rearrangement, Eq. (7) can be written as

$$i\partial\psi/\partial u = \Omega(u, x, y, p, \partial_x, \partial_y)\psi, \quad (8)$$

where Ω is a linear operator acting on the space of complex functions $\psi(u, x, y, p)$:

$$\Omega\psi = \{(\kappa^2\gamma - p^2\gamma^{33})\psi/\sqrt{\gamma^{03}} + [\gamma^{mn}(\psi/\sqrt{\gamma^{03}})_{,m}]_{,n} - 2ip\gamma^{m3}(\psi/\sqrt{\gamma^{03}})_{,m} - ip\gamma^{m3}_{,m}(\psi/\sqrt{\gamma^{03}})\}/2p\sqrt{\gamma^{03}}. \quad (9)$$

We are using here a mathematical technique borrowed from ordinary quantum mechanics, but at this stage the problem is completely classical.

It will now be shown that $\int \psi^* \Omega \psi dx dy$ is real (that is, Ω is a Hermitian operator) if ψ falls rapidly enough for large x and y . This result is obvious for the first two terms in the braces on the right hand side of (9). For the third term, we have, after integrating by parts,

$$\int \psi^* [(\gamma^{mn}\chi_{,m})_{,n}/\sqrt{\gamma^{03}}] dx dy = - \int \chi^*_{,n} \gamma^{mn} \chi_{,m} dx dy, \quad (10)$$

which is also real. Likewise, the last two terms together give

$$-(i/2) \int \chi^* (2\gamma^{m3}\chi_{,m} + \gamma^{m3}_{,m}\chi) dx dy = -(i/2) \int (\chi^* \gamma^{m3} \chi_{,m} - \chi^*_{,m} \gamma^{m3} \chi) dx dy, \quad (11)$$

where an integration by parts was performed on the second term on the left hand side. Again, the result is seen to be real.

It therefore follows from Eq. (8) that

$$(d/du) \int |\psi(u, x, y, p)|^2 dx dy = 0 \quad \forall p, \quad (12)$$

so that the expression $\int |\psi(u, x, y, p)|^2 dx dy$ is a constant of the motion (it has the same value on all the hypersurfaces $u = \text{const.}$). We thus have an infinity of conservation laws, one for each value of p .

We now turn our attention to the quantized version of the above theory. We assume that a quantum field Ψ satisfies the same Klein-Gordon equation as the classical field, for the given classical background $g_{\alpha\beta}$:

$$i\partial\Psi(u, x, y, p)/\partial u = \Omega(u, x, y, p, \partial_x, \partial_y)\Psi(u, x, y, p). \quad (13)$$

Note that Ψ is a quantum field operator, acting on the Hilbert space of quantum states, while Ω is a purely classical linear operator referring to the classical parameters u , x , y , and p . The problem is whether Eq. (13) generates a *unitary* transformation of the field operator Ψ .

This question is not trivial, because the present classical field theory does not have the usual canonical structure, and there is no guarantee that quantization is possible [9]. The term $\frac{1}{2}\gamma^{\alpha\beta}\phi_{,\alpha}\phi_{,\beta}$ in the Lagrangian density $\mathcal{L}$ produces a canonical momentum $\pi \equiv \partial\mathcal{L}/\partial\phi_{,0} = \gamma^{03}\phi_{,3}$. This relationship is a *constraint* restricting the values of π and ϕ on a constant x^0 hypersurface. Moreover, the Hamiltonian density, $\mathcal{H} \equiv \pi\phi_{,0} - \mathcal{L}$, does not depend on π at all! The analogy between Poisson brackets and commutators therefore fails, and it is not obvious that there is a unitary evolution law $i\partial\Psi(u, x, y, p)/\partial u = [\Psi(u, x, y, p), H]$, where H is a Hermitian operator acting on the quantum Hilbert space. This ought to be *proved*.

Since the classical ψ satisfied $\psi(u, x, y, -p) = \psi^*(u, x, y, p)$, we shall assume the same for the quantized field:

$$\Psi^\dagger(u, x, y, p) \equiv \Psi(u, x, y, -p), \quad (14)$$

where a dagger denotes the adjoint of an operator in the quantum Hilbert space (an asterisk denotes the complex conjugate of an ordinary number). We shall henceforth consider both Ψ and $\Psi^\dagger$ as independent variables, and on the other hand we restrict p to positive values [2,3]. We further postulate the equal- u commutation relations (whose consistency we shall have to check, since they cannot be derived from an analogy with classical Poisson brackets):

$$[\Psi(u, x', y', p'), \Psi(u, x'', y'', p'')] = 0, \quad (15)$$

$$[\Psi^\dagger(u, x', y', p'), \Psi^\dagger(u, x'', y'', p'')] = 0, \quad (16)$$

$$[\Psi(u, x', y', p'), \Psi^\dagger(u, x'', y'', p'')] = \delta(x' - x'')\delta(y' - y'')\delta(p' - p''), \quad (17)$$

where an arbitrary normalization constant on the right hand side of (17) has been ignored.

Finally, we define, in the quantum Hilbert space, a Hermitian operator H , which will play the role of a Hamiltonian, as we shall see:

$$H \equiv \int \Psi^\dagger(u, x, y, p) \Omega \Psi(u, x, y, p) dx dy dp. \quad (18)$$

We then indeed have

$$[\Psi(u, x', y', p'), H] = \Omega \Psi(u, x', y', p'), \quad (19)$$

and we can consistently postulate that both sides of (19) are equal to $i\partial\Psi(u, x', y', p')/\partial u$, in agreement with Eq. (13). This justifies the name ‘‘Hamiltonian’’ given to the operator H defined above. That operator generates a *unitary* dynamical evolution: the commutation relations (15–17) remain valid for all u .

This can also be seen in a more complicated but more general way, which will be useful later, because it does not rely on the existence of a Hamiltonian. We can directly differentiate Eq. (17) with respect to u , and use Eq. (13) and its adjoint,

$$i\partial\Psi^\dagger(u, x, y, p)/\partial u = \tilde{\Omega}(u, x, y, p, \partial_x, \partial_y) \Psi^\dagger(u, x, y, p), \quad (20)$$

where $\tilde{\Omega}(u, x, y, p, \partial_x, \partial_y) \equiv \Omega(u, x, y, -p, \partial_x, \partial_y)$, by virtue of (14). (We can also write this as $\tilde{\Omega} = -\Omega^*$.) Since Ω is an operator acting on functions of x, y , and p , and is local in these coordinates, it is straightforward (but tedious) to verify that

$$i(\partial/\partial u)[(\Psi)', (\Psi^\dagger)'] = [(\Omega\Psi)', (\Psi^\dagger)'] + [(\Psi)', (\tilde{\Omega}\Psi^\dagger)'] = 0, \quad (21)$$

where $(\Psi)'$ means $\Psi(u, x', y', p')$, and likewise for the other symbols. It is crucial in the verification of (21) that Ω be both local and Hermitian.

The commutation relations (15–17) show that the operators Ψ and $\Psi^\dagger$ behave as annihilation and creation operators, respectively. In a flat spacetime, it is possible to Fourier transform the coordinates x and y too, and it then becomes manifest that the transformed operators refer to particles of mass $\hbar\kappa/c$. [This will be explicitly shown later, see Eq. (29) and the ensuing discussion.]

Consider in particular the case of a “sandwich wave”: for $u < u_1$ and $u > u_2$, spacetime is flat ($g^{03} = -g^{11} = -g^{22} = 1$, other $g^{\alpha\beta}$ vanish). As we have just seen, there are *separate* linear field equations for Ψ and for $\Psi^\dagger$, even in the region $u_1 < u < u_2$ (this is related to the conservation of the classical momentum p_3). Nowhere is there any mixing of creation and annihilation operators. Therefore there is no Bogoliubov transformation. The number of particles is conserved, and in particular no particles emerge from the vacuum [4–6]. The meaning of the more general conservation laws that we found here is that *for each value of p_3 separately*, the number of particles with momentum p_3 is conserved.

Finally, we turn our attention to more general null-fronted metrics, where $g^{0m} \neq 0$. In that case, the classical field equations can be written as

$$A(i\partial\psi/\partial u) = B\psi, \quad (22)$$

where $\psi \equiv \sqrt{\gamma^{03}}\chi$ as before, and where A and B are linear operators acting on the complex function $\psi(u, x, y, p)$. Explicitly,

$$A = 2p + i[(\partial/\partial x^m)(\gamma^{0m}/\gamma^{03}) + (\gamma^{0m}/\gamma^{03})(\partial/\partial x^m)], \quad (23)$$

which is manifestly Hermitian. Then, with the exception of a subspace of functions satisfying $A\psi = 0$, there exists [10] a unique inverse operator, A^{-1} , and we can write

$$i\partial\psi/\partial u = (A^{-1}B)\psi. \quad (24)$$

Note that A^{-1} is not local (the inverse of a differential operator is an integral operator) but this is not the main issue. The difficulty is that $A^{-1}B$ is not, in general Hermitian. Consequently, if we attempt to have a quantized field operator Ψ which obeys an equation analogous to (24), the commutation relation (17) is in jeopardy.

Let us substitute $A^{-1}B$ for Ω in Eq. (21) (and likewise $\tilde{A}^{-1}\tilde{B}$ for $\tilde{\Omega}$, where a tilde on A or B means the substitution $p \rightarrow -p$), and let us multiply the result by $(A)'(\tilde{A})''$, to eliminate the nonlocality. We obtain an expression

$$[(B\Psi)', (\tilde{A}\Psi^\dagger)'] + [(A\Psi)', (\tilde{B}\Psi^\dagger)'], \quad (25)$$

and the question is whether this sum vanishes, so that (17) can hold for all u .

As a simple example, consider the case where the only nonvanishing components of $\gamma^{\alpha\beta}$ are $\gamma^{03} = -\gamma^{11} = -\gamma^{22} = 1$, and $\gamma^{01} = f(u)$. Let $g(u) \equiv df/du$. We have

$$A = 2p + 2i f(u) \partial/\partial x, \quad (26)$$

$$B = -(\partial^2/\partial x^2) - (\partial^2/\partial y^2) + \kappa^2 + g(u) \partial/\partial x. \quad (27)$$

Since the $\gamma^{\alpha\beta}$ do not depend on x or y , we can perform a Fourier transformation on these two coordinates, and define

$$a(u, \mathbf{k}, p) \equiv (2\pi)^{-1} \int e^{-i\mathbf{k}\cdot\mathbf{r}} \Psi(u, x, y, p) dx dy, \quad (28)$$

where $\mathbf{k} \equiv \{k_x, k_y\}$ and $\mathbf{r} = \{x, y\}$. We likewise define the adjoint operator $a^\dagger(u, \mathbf{k}, p)$. It is straightforward to show from Eq. (17) that

$$[a(u, \mathbf{k}', p'), a^\dagger(u, \mathbf{k}'', p'')] = \delta(\mathbf{k}' - \mathbf{k}'') \delta(p' - p''). \quad (29)$$

The Fourier transform of the field equation (22) is, with the values of A and B specified above,

$$2(p - f k_x) (i \partial a / \partial u) = (\mathbf{k}^2 + \kappa^2 + i g k_x) a, \quad (30)$$

and likewise for $a^\dagger$.

In the special case of flat space ($f = g = 0$), Eq. (30) reduces to the equation of motion (in the Heisenberg picture) for the annihilation operator, $a(u, \mathbf{k}, p)$, of a particle of mass $\hbar\kappa/c$. However, if $g \neq 0$, the situation is more intricate. Let us assume that the commutation relation (29) is valid at some “time” u_1 . At later times, we have, from (30),

$$a(u, \mathbf{k}, p) = a(u_1, \mathbf{k}, p) Z(u_1, u, \mathbf{k}, p), \quad (31)$$

where

$$Z(u_1, u, \mathbf{k}, p) = \exp \left(-i \int_{u_1}^u \frac{\mathbf{k}^2 + \kappa^2 + i g(u') k_x}{2[p - f(u') k_x]} du' \right). \quad (32)$$

Because of the ig term in the numerator, $|Z| \neq 1$ and Eq. (31) is not a unitary transformation. A straightforward calculation gives

$$[a(u, \mathbf{k}', p'), a^\dagger(u, \mathbf{k}'', p'')] = \delta(\mathbf{k}' - \mathbf{k}'') \delta(p' - p'') |Z(u_1, u, \mathbf{k}', p')|^2. \quad (33)$$

If this is transformed back into coordinate space, the result is not proportional to $\delta(x' - x'') \delta(y' - y'')$, but is

$$[(\Psi)', (\Psi^\dagger)''] = (2\pi)^{-1} \delta(p' - p'') \int |Z(u_1, u, \mathbf{k}, p)|^2 e^{i\mathbf{k}\cdot(\mathbf{r}' - \mathbf{r}'')} d\mathbf{k}. \quad (34)$$

This is a *nonlocal* function of $(x' - x'')$ and $(y' - y'')$. The commutation relation (17) is therefore incompatible with the field equation (13). The quantum field $\Psi(u, x, y, p)$ cannot be interpreted in terms of particle creation and annihilation operators, and the physical meaning of the theory is problematic. [In this particular case, there is a simple way out of the dilemma: returning to the usual time, $t = (u + v)/\sqrt{2}$. Surfaces of constant t have everywhere a time-like normal vector, and a Cauchy problem can be properly formulated. However, this may not be possible for more general metrics.]

Work by OL was supported by the Technion Graduate School. Work by AP was supported by the Gerard Swope Fund and the Fund for Encouragement of Research.

1. P. A. M. Dirac, *Rev. Mod. Phys.* **21**, 392 (1949).
2. H. Leutwyler, J. R. Klauder, and L. Streit, *Nuovo Cimento A* **66**, 536 (1970).
3. F. Coester, *Prog. Part. Nucl. Phys.* **29**, 1 (1992).
4. S. Deser, *J. Phys. A* **8**, 1972 (1975).
5. G. W. Gibbons, *Comm. Math. Phys.* **45**, 191 (1975).
6. J. Garriga and E. Verdaguer, *Phys. Rev. D* **43**, 391 (1991).
7. A. Peres, *Phys. Rev. Letters* **3**, 571 (1959).
8. A. Peres, *Phys. Rev.* **118**, 1105 (1960).
9. S. A. Hojman and L. C. Shepley, *J. Math. Phys.* **32**, 142 (1991).
10. F. Riesz F and B. Sz.-Nagy, *Functional Analysis*, Ungar, New York (1955) p. 152.

QUANTUM THEORY: CONCEPTS AND METHODS

by Asher Peres

446 + xiv pages

ISBN 0-7923-2549-4

in USA and Canada:

Kluwer Academic Publishers
P. O. Box 358
Accord Station
Hingham, MA 02018-0358

Phone 617-871-6600
Fax 617-871-6528

in all other countries:

Kluwer Academic Publishers
P. O. Box 322
3300 AH Dordrecht
The Netherlands

Phone (0)78 - 524400
Fax (0)78 - 524474

TABLE OF CONTENTS

Chapter 1: Introduction to Quantum Physics

1-1. The downfall of classical concepts	3
1-2. The rise of randomness	5
1-3. Polarized photons	7
1-4. Introducing the quantum language	9
1-5. What is a measurement?	14
1-6. Historical remarks	18
1-7. Bibliography	21

Chapter 2: Quantum Tests

2-1. What is a quantum system?	24
2-2. Repeatable tests	27
2-3. Maximal quantum tests	29
2-4. Consecutive tests	33
2-5. The principle of interference	36
2-6. Transition amplitudes	39
2-7. Appendix: Bayes's rule of statistical inference	45
2-8. Bibliography	47

Chapter 3: Complex Vector Space

3-1. The superposition principle	48
3-2. Metric properties	51
3-3. Quantum expectation rule	54
3-4. Physical implementation	57
3-5. Determination of a quantum state	58
3-6. Measurements and observables	62
3-7. Further algebraic properties	67
3-8. Quantum mixtures	72
3-9. Appendix: Dirac's notation	77
3-10. Bibliography	78

Chapter 4: Continuous Variables

4-1. Hilbert space	79
4-2. Linear operators	84
4-3. Commutators and uncertainty relations	89
4-4. Truncated Hilbert space	95
4-5. Spectral theory	99
4-6. Classification of spectra	103
4-7. Appendix: Generalized functions	106
4-8. Bibliography	112

Chapter 5: Composite Systems

5-1. Quantum correlations	115
5-2. Incomplete tests and partial traces	121
5-3. The Schmidt decomposition	123
5-4. Indistinguishable particles	126
5-5. Parastatistics	131
5-6. Fock space	137
5-7. Second quantization	142
5-8. Bibliography	147

Chapter 6: Bell's Theorem

6-1. The dilemma of Einstein, Podolsky, and Rosen	148
6-2. Cryptodeterminism	155
6-3. Bell's inequalities	160
6-4. Some fundamental issues	167
6-5. Other quantum inequalities	173
6-6. Higher spins	179
6-7. Bibliography	185

Chapter 7: Contextuality

7-1. Nonlocality versus contextuality	187
7-2. Gleason's theorem	190
7-3. The Kochen-Specker theorem	196
7-4. Experimental and logical aspects of inseparability	202
7-5. Appendix: Computer test for Kochen-Specker contradiction	209
7-6. Bibliography	211

Chapter 8: Spacetime Symmetries

8-1. What is a symmetry?	215
8-2. Wigner's theorem	217
8-3. Continuous transformations	220
8-4. The momentum operator	225
8-5. The Euclidean group	229
8-6. Quantum dynamics	237
8-7. Heisenberg and Dirac pictures	242
8-8. Galilean invariance	245
8-9. Relativistic invariance	249
8-10. Forms of relativistic dynamics	254
8-11. Space reflection and time reversal	257
8-12. Bibliography	259

Chapter 9: Information and Thermodynamics

9-1. Entropy	260
9-2. Thermodynamic equilibrium	266
9-3. Ideal quantum gas	270
9-4. Some impossible processes	275
9-5. Generalized quantum tests	279
9-6. Neumark's theorem	285
9-7. The limits of objectivity	289
9-8. Quantum cryptography and teleportation	293
9-9. Bibliography	296

Chapter 10: Semiclassical Methods

10-1. The correspondence principle	298
10-2. Motion and distortion of wave packets	302
10-3. Classical action	307
10-4. Quantum mechanics in phase space	312
10-5. Koopman's theorem	317
10-6. Compact spaces	319
10-7. Coherent states	323

10-8. Bibliography	330
Chapter 11: Chaos and Irreversibility	
11-1. Discrete maps	332
11-2. Irreversibility in classical physics	341
11-3. Quantum aspects of classical chaos	347
11-4. Quantum maps	351
11-5. Chaotic quantum motion	353
11-6. Evolution of pure states into mixtures	369
11-7. Appendix: PostScript code for a map	370
11-8. Bibliography	371
Chapter 12: The Measuring Process	
12-1. The ambivalent observer	373
12-2. Classical measurement theory	378
12-3. Estimation of a static parameter	385
12-4. Time-dependent signals	387
12-5. Quantum Zeno effect	392
12-6. Measurements of finite duration	400
12-7. The measurement of time	405
12-8. Time and energy complementarity	413
12-9. Incompatible observables	417
12-10. Approximate reality	423
12-11. Bibliography	428
Author Index	430
Subject Index	435